


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2007, The Johns Hopkins University and James Dick. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.



JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Overview of Microbiology

James D. Dick, PhD
Johns Hopkins University

- Director, Bacteriology Section,
Division of Medical Microbiology
- Associate Professor of Pathology,
Molecular Microbiology and Immunology
- Researches biochemical and molecular
techniques for the detection and identi-
fication of bacterial pathogens and anti-
biotic resistance mechanisms



JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Section A

Overview of Microbiology

Classification of Infectious Disease

- Clinical
 - Major clinical manifestation
- Epidemiological
 - Transmission/reservoir
- Microbiological
 - Causative agent

Microbiological Classification of Infectious Disease

- Viral
- Bacterial
- Fungal
- Parasitic
- Prions


Diagnosis of Infectious Diseases

- Infection versus colonization
- Disease versus exposure
- Prevalence versus incidence

Comparison of Eukaryotic and Prokaryotic Cells


Characteristic	Eukaryotes	Prokaryotes
Form	Multicellular	Single cells
Nucleus	Nuclear membrane	DNA in contact with cytoplasm
Organelles	Membrane-bound organelles present	No organelles
Sterols	Always	Only in <i>Mycoplasma</i>
Ribosomes	80s = 40s + 60s	70s = 30s + 50s
Cell wall	Absent or cellulose/chitin	Peptidoglycan
Mitosis	Yes	No

Size Comparisons of Microorganisms



Bacterial Cell Structure

Bacterial Cell Structure



Bacterial Classification

- Gram-positive
- Gram-negative
- Acid-fast

Bacterial Shapes

- Cocci-round or spherical cells
- Bacilli-rod-shaped cells
- Curved, spiral forms

Biological Stains


- Bacteria—gram stain
- Fungi—KOH, lactophenol blue, India ink, silver stains in tissue
- Mycobacteria—acid fast stains
- Parasites—trichrome stain, Wright's stain
- Viruses—antibody conjugated dyes

Spherical, Rod-Shaped, and Spiral


Cocci
(spherical)



Staphylococcus aureus



Streptococcus pneumoniae



Streptococcus pyogenes

Bacilli
(rods)



Bacillus anthracis




Haemophilus influenzae

Curved or
spiral



Vibrio cholerae




Borrelia burgdorferi

Gram-positive


Gram-negative

Bacterial Cell Walls

Gram +



Gram -



Pappenheim Stain of Streptococcus pyogenes



Public Domain


Dark Field Microscopy of Syphilis









Classification of Viruses

Family	Example	Genome Size, Kilobases of Kilobase Pairs	Envelope
RNA Viruses			
Single-stranded			
Picornaviridae	Poliovirus	7.2-8.4	No
Togaviridae	Rubella virus	12	Yes
Flaviviridae	Yellow fever virus	10	Yes
Coronaviridae	SARS	16-21	Yes
Rhabdoviridae	Rabies virus	13-16	Yes
Paramyxoviridae	Measles virus	16-20	Yes
Orthomyxoviridae	Influenza virus	14	Yes
Bunyaviridae	California encephalitis virus	13-21	Yes
Arenaviridae	Lymphocytic choriomeningitis virus	10-14	Yes
Retroviridae	HIV	319	Yes
Double-stranded			
Reoviridae	Rotaviruses	16-27	No
DNA Viruses			
Single-stranded			
Parvoviridae	Human parvovirus B-19	5	No
Partially double-stranded			
Hepadnaviridae	Hepatitis B	3	Yes
Double-stranded			
Polyomaviridae	JC virus	8	No
Adenoviridae	Human adenovirus	36-38	No
Herpesviridae	Herpes simplex virus	120-220	Yes
Poxviridae	Vaccinia, smallpox	130-280	Yes


Nonenveloped

Double-stranded DNA		Double-stranded RNA	Single-stranded RNA		Single-stranded DNA
					
Papovaviridae (Papilloma)	Adenoviridae (Adeno)	Reoviridae (Reo)	Calicivirus (Norwalk?)	Picornaviridae (Polio)	Parvoviridae (Erythema infectiosum)

Enveloped


Double-stranded DNA		Single-stranded RNA			
					
Poxviridae (Vaccinia)	Herpesviridae (Herpes simplex)	Paramyxoviridae (Measles)	Orthomyxoviridae (Influenza)	Rhabdoviridae (Rabies)	Retroviridae (Human immunodeficiency)
					
Hepadnaviridae (Hepatitis B)	Flaviviridae (St. Louis encephalitis)	Arenaviridae (Lymphocytic choriomeningitis)	Coronaviridae (Coronavirus)	Bunyaviridae (Hantavirus)	Togaviridae (Rubella)

Electron Micrograph of Viral Particle (Influenza)



Public Domain

Fluorescent Antibody Stain (Meningoencephalitis)



Characteristics of Fungi

- Pathogenic fungi have two forms: yeasts (unicellular) and molds (multicellular)
- Some fungi are dimorphic
- Molds grow as filamentous, branching strands of connected cells known as hyphae

Characteristics of Fungi

- Fungi classified by type/method of reproduction
- Asexual—conidia; sexual—spores
- “Micro” or “macro” refers to size of spores


Grouping of Fungi of Medical Importance

- Superficial mycoses—outermost layers of skin and hair
- Cutaneous mycoses—epidermis
- Subcutaneous mycoses—dermis and subcutaneous tissues
- Systemic mycoses—internal organ systems

Fungi of Medical Importance

Fungus	Classification	Disease
<i>Malassezia furfur</i>	Yeast	Superficial mycoses
<i>Trichophyton rubrum</i>	Filamentous	Tinea, cutaneous mycoses
<i>Microsporum audouinii</i>	Filamentous	Tinea, cutaneous mycoses
<i>Epidermophyton floccosum</i>	Filamentous	Tinea, cutaneous mycoses
<i>Candida albicans</i>	Yeast	Mucocutaneous and systemic mycoses
<i>Sporothrix schenckii</i>	Dimorphic	Subcutaneous mycoses
<i>Histoplasma capsulatum</i>	Dimorphic	Systemic mycoses, histoplasmosis
<i>Blastomyces dermatitidis</i>	Dimorphic	Systemic mycoses, blastomycosis
<i>Coccidioides immitis</i>	Dimorphic	Systemic mycoses, coccidioidomycosis
<i>Paracoccidioides brasiliensis</i>	Dimorphic	Systemic mycoses, paracoccidioidomycosis
<i>Penicillium marneffeii</i>	Dimorphic	Systemic mycoses, penicilliosis
<i>Cryptococcus neoformans</i>	Yeast	Systemic mycoses, opportunistic cryptococcosis
<i>Candida</i> species	Yeast	Opportunistic infections
<i>Aspergillus fumigatus</i>	Filamentous	Opportunistic infections
<i>Aspergillus flavus</i>	Filamentous	Opportunistic infections
<i>Fusarium</i> species	Filamentous	Opportunistic infections
<i>Rhizopus</i> species	Filamentous	Opportunistic infections
<i>Mucor</i> species	Filamentous	Opportunistic infections
<i>Absidia</i> species	Filamentous	Opportunistic infections
<i>Pseudallesheria</i> species	Filamentous	Opportunistic infections
<i>Pneumocystis carinii</i>	Dimorphic, cysts	Opportunistic infections

Histoplasma capsulatum



Public Domain

- Protozoa
 - Amebiasis
 - Leishmaniasis
 - Trypanosomiasis
 - Malaria
- Helminths—adult stage in worm

Hookworm in stool





JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Section B

Diagnostic Microbiology

- Microscopy
- Culture
- Immunology
- Molecular methods

- Microscopy
 - Light microscopy
 - Fluorescence microscopy
 - Electron microscopy


Bacteria

- Staining characteristics
 - Gram negative/gram positive
- Shape
 - Cocci-round or spherical cells
 - Bacilli-rod-shaped cells
 - Curved, spiral forms
 - Coccobacilli
 - Pleomorphic cells
- Size

- Sensitivity
 - Relatively poor, one bacterial cell per high-powered field is equivalent to 100,000–1,000,000 bacteria per ml
 - This can be improved through fluorescence
- Specificity
 - Poor for bacteria, high for microorganisms exhibiting distinctive morphology—filamentous fungi, parasites, viruses by EM

- Culture
 - Defined versus undefined media
 - Enrichment media
 - Selective media
 - Specialized media

Blood Agar (*Corynebacterium haemolyticum*)



MacConkey's Agar (Proteus Vulgaris)




Public Domain

- Sensitivity—usually considered the gold standard
- Specificity—excellent when used in conjunction with phenotypic, immunologic, and molecular techniques


- Testing for specific microbial antigens
 - Direct detection from clinical specimens
 - Characterization of a cultured organism
- Testing for antibody to specific microbial antigens
 - Detection of a particular isotope, usually IgM or IgG
 - IgA and IgE not usually used

- Complement fixation
- Agglutination assays
- Neutralization/hemagglutination assays
- Enzyme immunoassays (EIAs, ELISAs)
- Radioimmunoassays (RIA)
- Fluorescent antibody techniques

Fluorescent Antibody Stain (Neisseria gonorrhoeae)




Microbiology Tools for the Epidemiologist



- Nucleic acid probes
- Signal amplification methods
 - PCR
 - RT-PCR
 - Nested PCR
 - Multiplex PCR

Sensitivity Comparisons of Molecular Methods




- Detection of fastidious, slow-growing, or nonculturable organisms as well as antibiotic resistance genes and phenotypically difficult organisms


Hybridization Techniques

- Solid-phase
- Liquid phase
- *In situ*

Polymerase Chain Reaction (PCR)



Gel and Hybridizations of PCR Reactions



Limitations of Molecular Technology

- False positive results
 - Contamination
- False negative results
 - Presence of inhibitors
- Narrow spectrum of detection
 - Detection of a single pathogen
- Cost